

THE CYLINDRICAL CONTACT HOMOLOGY OF UNIVERSALLY TIGHT SUTURED CONTACT SOLID TORI

ROMAN GOLOVKO

ABSTRACT. We calculate the sutured version of cylindrical contact homology of a sutured contact solid torus $(S^1 \times D^2, \Gamma, \xi)$, where Γ is arbitrary and ξ is a universally tight contact structure.

1. INTRODUCTION

The cylindrical contact homology of a (closed) contact manifold was introduced by Eliashberg and Hofer and is the simplest version of the symplectic field theory of Eliashberg, Givental and Hofer [6]. It is the homology of a differential graded module whose differential counts genus zero holomorphic curves in the symplectization with one positive puncture and one negative puncture.

A natural condition to impose on a compact, oriented contact $(2m + 1)$ -manifold (M, ξ) with boundary is to require that ∂M be *convex*, i.e., there is a contact vector field X transverse to ∂M . To a transverse contact vector field X we can associate the *dividing set* $\Gamma = \Gamma_X \subset \partial M$, namely the set of points $x \in \partial M$ such that $X(x) \in \xi(x)$. By the contact condition, $(\Gamma, \xi \cap T\Gamma)$ is a $(2m - 1)$ -dimensional contact submanifold of (M, ξ) ; the isotopy class of $(\Gamma, \xi \cap T\Gamma)$ is independent of the choice of X . We will denote by (M, Γ, ξ) the contact manifold (M, ξ) with convex boundary and dividing set $\Gamma = \Gamma_X \subset \partial M$ with respect to some transverse contact vector field X . Note that the actual boundary condition we need is slightly different and is called a *sutured boundary condition*. (In the early 1980's, Gabai developed the theory of sutured manifolds [7], which became a powerful tool in studying 3-manifolds with boundary.) For the moment we write (M, Γ, ξ) to indicate either the convex boundary condition or the sutured boundary condition.

It turns out that there is a way to generalize cylindrical contact homology to sutured manifolds. This is possible by imposing a certain convexity condition on the contact form. This construction is described in the paper of Colin, Ghiggini, Honda and Hutchings [3] and will be summarized in Section 2.

In this paper, we construct a sutured contact solid torus with $2n$ parallel sutures of slope $\frac{k}{l}$ using the gluing method of Colin, Ghiggini, Honda and Hutchings [3], and calculate the sutured cylindrical contact homology of it. Here $n \in \mathbb{N}$, $(k, l) = 1$ and $|k| > l > 0$. In order to define the slope, we choose an oriented identification $\partial(S^1 \times D^2) \simeq T^2 = (\mathbb{R}/\mathbb{Z})^2$ as follows: map $\{pt\} \times \partial D^2$ (the meridian) to $(1, 0)$ (slope is 0) and $S^1 \times \{pt\}$ (a longitude) to $(0, 1)$.

This calculation, together with the calculation of the sutured cylindrical contact homology of the sutured contact solid torus with $2n$ parallel longitudinal sutures, where $n \geq 2$, that has been done in [8], finishes the calculation of the cylindrical contact homology of $(S^1 \times D^2, \Gamma, \xi)$, where Γ is arbitrary and ξ is a universally tight contact structure.

2000 *Mathematics Subject Classification.* Primary 53D42; Secondary 57M50, 53D10.

Key words and phrases. sutured manifolds, contact homology.

Our goal is to prove the following theorem:

Theorem 1.1. *Let $(S^1 \times D^2, \Gamma)$ be a sutured manifold, where Γ is a set of $2n$ parallel closed curves of slope $\frac{k}{l}$, where $(k, l) = 1$, $|k| > l > 0$ and $n \in \mathbb{N}$. Then there is a contact form α which makes $(S^1 \times D^2, \Gamma, \alpha)$ a sutured contact manifold with a universally tight contact structure $\xi = \ker \alpha$, $HC^{cyl}(S^1 \times D^2, \Gamma, \alpha)$ is defined, is independent of the contact form α for $\xi = \ker \alpha$ and the almost complex structure J and*

$$HC^{cyl, h}(S^1 \times D^2, \Gamma, \xi) \simeq \begin{cases} \mathbb{Q}, & \text{for } k \nmid h > 0; \\ \mathbb{Q}^{n-1}, & \text{for } k \mid h > 0; \\ 0, & \text{otherwise.} \end{cases}$$

Here h corresponds to the homological grading.

ACKNOWLEDGEMENTS

The author is deeply grateful to Ko Honda for his guidance, help and support. He also thanks Dmytro Chebotarov, Oliver Fabert, Paolo Ghiggini, Jian He, Michael Hutchings and Mark McLean for helpful suggestions and interest in his work. In addition, the author is extremely grateful to Andrew Cotton-Clay for his critical comments on the first version of the paper. Finally, the author thanks the Mathematical Sciences Research Institute and the organizers of the ‘‘Symplectic and Contact Geometry and Topology’’ program for their hospitality.

2. BACKGROUND

The goal of this section is to review definitions of sutured contact manifold and the relative version of cylindrical contact homology. This section can be considered as a summary of [3].

2.1. Review of sutured contact manifolds. In this section we recall some definitions and describe some constructions from [3]. We first start with the notion of a Liouville manifold.

Definition 2.1. A *Liouville manifold* (often also called a Liouville domain) is a pair (W, β) consisting of a compact, oriented $2n$ -dimensional manifold W with boundary and a 1-form β on W , where $\omega = d\beta$ is a positive symplectic form on W and the *Liouville vector field* Y given by $i_Y(\omega) = \beta$ is positively transverse to ∂W . It follows that the 1-form $\beta_0 = \beta|_{\partial W}$ (this notation means β pulled back to ∂W) is a positive contact form with kernel ζ .

We now recall the definition of a sutured contact manifold.

Definition 2.2. A compact oriented $2n + 1$ -dimensional manifold M with boundary and corners is a *sutured contact manifold* if it comes with an oriented, not necessarily connected submanifold $\Gamma \subset \partial M$ of dimension $2n - 1$ (called the *suture*), together with a neighborhood $U(\Gamma) = [-1, 0] \times [-1, 1] \times \Gamma$ of $\Gamma = \{0\} \times \{0\} \times \Gamma$ in M , with coordinates $(\tau, t) \in [-1, 0] \times [-1, 1]$, such that the following holds:

- (1) $U \cap \partial M = (\{0\} \times [-1, 1] \times \Gamma) \cup ([-1, 0] \times \{-1\} \times \Gamma) \cup ([-1, 0] \times \{1\} \times \Gamma)$;
- (2) $\partial M \setminus (\{0\} \times (-1, 1) \times \Gamma) = R_-(\Gamma) \sqcup R_+(\Gamma)$, where the orientation of ∂M agrees with that of $R_+(\Gamma)$ and is opposite that of $R_-(\Gamma)$ and the orientation of Γ agrees with the boundary orientation of $R_+(\Gamma)$;
- (3) the corners of M are precisely $\{0\} \times \{\pm 1\} \times \Gamma$.

In addition, M is equipped with a contact structure ξ , which is given by the kernel of a positive contact 1-form α such that:

- (i) $(R_{\pm}(\Gamma), \beta_{\pm} = \alpha|_{R_{\pm}(\Gamma)})$ is a Liouville manifold;
- (ii) $\alpha = Cdt + \beta$ inside $U(\Gamma)$, where $C > 0$ and β is independent of t and does not have a dt -term;
- (iii) $\partial_{\tau} = Y_{\pm}$, where $Y_{\pm}$ is a Liouville vector field for $\beta_{\pm}$.

Such a contact form α is said to be *adapted* to $(M, \Gamma, U(\Gamma))$.

Here we briefly describe the way to glue sutured contact manifolds. This procedure was first described by Colin and Honda in [4] and then generalized by Colin, Ghiggini, Honda and Hutchings in [3].

Let $(M', \Gamma', U(\Gamma'), \xi')$ be a sutured contact 3-manifold with an adapted contact form α' . We denote by π the projection along ∂_t defined on $U(\Gamma')$.

Take 2-dimensional submanifolds $P_{\pm} \subset R_{\pm}(\Gamma')$ such that $\partial P_{\pm}$ is the union of $(\partial P_{\pm})_{\partial} \subset \partial R_{\pm}(\Gamma')$, $(\partial P_{\pm})_{int} \subset int(R_{\pm}(\Gamma'))$ and $\partial P_{\pm}$ is positively transversal to the Liouville vector field $Y'_{\pm}$ on $R_{\pm}(\Gamma')$. Whenever we refer to $(\partial P_{\pm})_{int}$ and $(\partial P_{\pm})_{\partial}$, we assume that closures are taken as appropriate. Moreover we make the assumption that $\pi((\partial P_{-})_{\partial}) \cap \pi(\partial P_{+})_{\partial} = \emptyset$.

Let φ be a diffeomorphism which sends $(P_{+}, \beta'_{+}|_{P_{+}})$ to $(P_{-}, \beta'_{-}|_{P_{-}})$ and takes $(\partial P_{+})_{int}$ to $(\partial P_{-})_{\partial}$ and $(\partial P_{+})_{\partial}$ to $(\partial P_{-})_{int}$. Note that, since $\dim M = 3$, we only need $\beta'_{+}|_{P_{+}}$ and $\varphi^{*}(\beta'_{-}|_{P_{-}})$ to match up on ∂P_{+} , since we can linearly interpolate between primitives of positive area forms on a surface.

Topologically, we construct the sutured manifold (M, Γ) from (M', Γ') and the gluing data (P_{+}, P_{-}, φ) as follows: Let $M = M' / \sim$, where

- $x \sim \varphi(x)$ for all $x \in P_{+}$;
- $x \sim x'$ if $x, x' \in \pi^{-1}(\Gamma')$ and $\pi(x) = \pi(x') \in \Gamma'$.

Then

$$R_{\pm}(\Gamma) = \frac{\overline{R_{\pm}(\Gamma') \setminus P_{\pm}}}{(\partial P_{\pm})_{int}} \sim \pi_{\pm}((\partial P_{\mp})_{\partial})$$

and

$$\Gamma = \frac{\overline{\Gamma' \setminus \pi(\partial P_{+} \sqcup \partial P_{-})}}{\pi((\partial P_{+})_{int} \cap (\partial P_{+})_{\partial})} \sim \pi((\partial P_{-})_{int} \cap (\partial P_{-})_{\partial}).$$

For the detailed description of the gluing procedure we refer to [3].

Finally, we describe the way to complete sutured contact manifold (M, α) to a noncompact contact manifold (M^{*}, α^{*}) . This construction was first described in [3].

Let $(M, \Gamma, U(\Gamma), \xi)$ be a sutured contact manifold with an adapted contact form α . The form α is then given by $Cdt + \beta_{\pm}$ on $[1 - \varepsilon, 1] \times R_{+}(\Gamma)$ and $[-1, -1 + \varepsilon] \times R_{-}(\Gamma)$ of $R_{+}(\Gamma) = \{1\} \times R_{+}(\Gamma)$ and $R_{-}(\Gamma) = \{-1\} \times R_{-}(\Gamma)$, where $t \in [-1, -1 + \varepsilon] \cup [1 - \varepsilon, 1]$ extends the t -coordinate on U . On U , $\alpha = Cdt + \beta$, $\beta = \beta_{+} = \beta_{-}$ and ∂_{τ} is a Liouville vector field Y for β . We first extend α to $[1, \infty) \times R_{+}(\Gamma)$ and $(-\infty, -1] \times R_{-}(\Gamma)$ by taking $Cdt + \beta_{\pm}$ as appropriate. The boundary of this new manifold is $\{0\} \times \mathbb{R} \times \Gamma$. Notice that since $\partial_{\tau} = Y$, the form $d\beta|_{[-1, 0] \times \{t\} \times \Gamma}$ is the symplectization of $\beta|_{\{0\} \times \{t\} \times \Gamma}$ in the positive τ -direction. We glue $[0, \infty) \times \mathbb{R} \times \Gamma$ with the form $Cdt + e^{\tau} \beta_0$, where β_0 is the pullback of β to $\{0\} \times \{t\} \times \Gamma$.

We denote by M^* the noncompact extension of M described above and by α^* the extension of α to M^* .

2.2. Review of cylindrical contact homology. Let $(M, \Gamma, U(\Gamma), \xi)$ be a sutured contact manifold with an adapted contact form α and (M^*, α^*) be its completion.

The *Reeb vector field* R_{α^*} that is associated to a contact form α^* is given by $d\alpha^*(R_{\alpha^*}, \cdot) = 0$ and $\alpha^*(R_{\alpha^*}) = 1$.

We assume that R_{α^*} is *nondegenerate*, i.e., the first return map along each (not necessarily simple) periodic orbit does not have 1 as an eigenvalue. Observe that nondegeneracy can always be achieved by a small perturbation.

Remark 2.3. Note that every periodic orbit of R_{α^*} lies in M . Hence, the set of periodic Reeb orbits of R_{α^*} coincides with the set of periodic Reeb orbits of R_α .

A Reeb orbit γ is called *elliptic* or *positive* (respectively *negative*) *hyperbolic* if the eigenvalues of P_γ are on the unit circle or the positive (resp. negative) real line respectively.

If τ is a trivialization of ξ over γ , we can then define the Conley-Zehnder index. In 3-dimensional situation, we can explicitly describe the Conley-Zehnder index and its behavior under multiple covers as follows:

Proposition 2.4 ([10]). *If γ is elliptic, then there is an irrational number $\phi \in \mathbb{R}$ such that P_γ is conjugate in $SL_2(\mathbb{R})$ to a rotation by angle $2\pi\phi$ and*

$$\mu_\tau(\gamma^k) = 2\lfloor k\phi \rfloor + 1,$$

where $2\pi\phi$ is the total rotation angle with respect to τ of the linearized flow around the orbit.

If γ is positive (respectively negative) hyperbolic, then there is an even (respectively odd) integer r such that the linearized flow around the orbit rotates the eigenspaces of P_γ by angle πr with respect to τ and

$$\mu_\tau(\gamma^k) = kr.$$

A closed orbit of R_{α^*} is said to be *good* if it does not cover a simple orbit γ an even number of times, where the first return map $\xi_{\gamma_0} \rightarrow \xi_{\gamma_T}$ has an odd number of eigenvalues in the interval $(-1, 0)$. Here T is the period of the orbit γ . An orbit that is not good is called *bad*.

We now recall the notion of an almost complex structure on $\mathbb{R} \times M^*$ that is tailored to (M^*, α^*) .

Let (W, β) be a Liouville manifold and ζ be the contact structure given on ∂W by $\ker(\beta_0)$, where $\beta_0 = \beta|_{\partial W}$. In addition, let $(\widehat{W}, \widehat{\beta})$ be the completion of (W, β) , i.e., $\widehat{W} = W \cup ([0, \infty) \times \partial W)$ and $\widehat{\beta}|_{[0, \infty) \times \partial W} = e^\tau \beta_0$, where τ is the $[0, \infty)$ -coordinate. An almost complex structure J_0 on $\widehat{W}$ is *$\widehat{\beta}$ -adapted* if J_0 is β_0 -adapted on $[0, \infty) \times \partial W$; and $d\beta(v, J_0 v) > 0$ for all nonzero tangent vectors v on W .

Definition 2.5. Let $(M, \Gamma, U(\Gamma), \xi)$ be a sutured contact manifold, α be an adapted contact form and (M^*, α^*) be its completion. We say that an almost complex structure J on $\mathbb{R} \times M^*$ is *tailored* to (M^*, α^*) if the following hold:

- (1) J is α^* -adapted, i.e, J is $\mathbb{R}$ -invariant, $J(\xi) = \xi$, $d\alpha(v, Jv) > 0$ for nonzero $v \in \xi$ and $J(\partial_s) = R_{\alpha^*}$, where s denotes the $\mathbb{R}$ -coordinate;
- (2) J is ∂_t -invariant in a neighborhood of $M^* \setminus \text{int}(M)$;

- (3) The projection of J to $T\widehat{R_\pm(\Gamma)}$ is a $\widehat{\beta}_\pm$ -adapted almost complex structure J_0 on the completion $(\widehat{R_+(\Gamma)}, \widehat{\beta}_+) \sqcup (\widehat{R_-(\Gamma)}, \widehat{\beta}_-)$ of the Liouville manifold $(R_+(\Gamma), \beta_+) \sqcup (R_-(\Gamma), \beta_-)$. Moreover, the flow of ∂_t identifies $J_0|_{\widehat{R_+(\Gamma)} \setminus R_+(\Gamma)}$ and $J_0|_{\widehat{R_-(\Gamma)} \setminus R_-(\Gamma)}$.

Given a sutured contact manifold $(M, \Gamma, U(\Gamma), \alpha)$ and an α^* -adapted almost complex structure J , we define the *sutured cylindrical contact homology group* $HC^{cyl}(M, \Gamma, \alpha, J)$ to be the cylindrical contact homology of (M^*, α^*, J) . The cylindrical contact homology chain complex $C(\alpha, J)$ is a $\mathbb{Q}$ -module freely generated by all good Reeb orbits, where the grading $|\cdot|$ and the boundary map ∂ are defined as in [1] with respect to the α^* -adapted almost complex structure J . The homology of $C(\alpha, J)$ is the sutured cylindrical contact homology group $HC^{cyl}(M, \Gamma, \alpha, J)$.

For our calculations we need the following construction of a “global” symplectic trivialization described in [1]. Assume that all the Reeb orbits of R_α are good. Let us now choose trivializations $\tau(\gamma)$ consistently for all Reeb orbits γ . Assume that $H_1(M; \mathbb{Z})$ is a free module. We pick representatives $C_1, \dots, C_s$ in $H_1(M; \mathbb{Z})$ for a basis of $H_1(M; \mathbb{Z})$, together with a trivialization of ξ along each representative C_i , $i = 1, \dots, s$. Now for a Reeb orbit γ , we distinguish the following cases:

- (1) $[\gamma] = 0 \in H_1(M; \mathbb{Z})$. Choose a spanning surface S_γ and use it to trivialize ξ along γ .
- (2) $0 \neq [\gamma] \in H_1(M; \mathbb{Z})$. We choose a surface S_γ realizing a homology between γ and a linear combination of the representatives C_i , $i = 1, \dots, s$. We then use S_γ to extend the chosen trivializations of ξ along the C_i , $i = 1, \dots, s$ to γ .

We denote the obtained trivialization by τ .

To a J -holomorphic curve in $\mathcal{M}^J(\gamma; \gamma')$, we can glue the chosen surfaces S_γ and $S_{\gamma'}$ and obtain a closed surface in M . Let $A \in H_2(M; \mathbb{Z})$ be its homology class; we can use it to decorate the corresponding connected component $\mathcal{M}_A^J(\gamma; \gamma')$ of the moduli space. Using τ we can write

$$(2.2.1) \quad \text{ind}(u) = |\gamma| - |\gamma'| + 2\langle c_1(\xi), A \rangle$$

for $u \in \mathcal{M}_A^J(\gamma; \gamma')$, where $|\gamma|$ is the *Conley-Zehnder grading* of γ defined by

$$(2.2.2) \quad |\gamma| := \mu_\tau(\gamma) - 1.$$

We will use Formulas 2.2.1 and 2.2.2 for our calculations.

In addition, we will need the following fact, which is a consequence of Lemma 5.4 in [2]:

Fact 2.6. *Let (M, α) be a closed, oriented contact manifold with nondegenerate Reeb orbits and*

$$u = (a, f) : (\dot{S}, j) \rightarrow (\mathbb{R} \times M, J)$$

be a J -holomorphic curve in $\mathcal{M}^J(\gamma; \gamma')$, where γ and γ' are good Reeb orbits, J is an α -adapted almost complex structure on $\mathbb{R} \times M$ and $\mathcal{M}^J(\gamma; \gamma')$ is a moduli space of J -holomorphic curves that we consider in cylindrical contact homology. Then the following inequality holds:

$$\mathcal{A}(\gamma) \stackrel{\text{def}}{=} \int_\gamma \alpha \geq \int_{\gamma'} \alpha \stackrel{\text{def}}{=} \mathcal{A}(\gamma')$$

with equality if and only if $\gamma = \gamma'$ and in this case the moduli space consists of a single element $\mathbb{R} \times \gamma$.

Now we recall the following theorem:

Theorem 2.7 ([1]). *Let (M, α) be a closed, oriented contact manifold with nondegenerate Reeb orbits. Let $C_m^h(M, \alpha)$ be the cylindrical contact homology complex, where h is a homotopy class of Reeb orbits and m corresponds to the Conley-Zehnder grading. If $C_k^0(M, \alpha) = 0$ for $k = -1, 0, 1$, then for every free homotopy class h*

- (1) $\partial^2 = 0$;
- (2) $H(C_*^h(M, \alpha), \partial)$ is independent of the contact form α for ξ , the almost complex structure J and the choice of perturbation for the moduli spaces.

When M is closed and $\mathbb{R} \times M$ is 4-dimensional, the following transversality result has been proven by Momin, see Proposition 2.10 in [11]:

Theorem 2.8 ([11]). *Let $u \in \mathcal{M}^J(\gamma; \gamma')$ be such that $\text{ind}(u) = 1$. Then the linearization of the Cauchy-Riemann operator is surjective at u .*

Remark 2.9. Observe that Theorem 2.8 does not require J to be generic.

Remark 2.10. Note that Theorem 2.8 can be considered as a consequence of the automatic transversality result of Wendl, see Theorem 0.1 in [12].

Finally, we recall the following result of Colin, Ghiggini, Honda and Hutchings from [3]:

Theorem 2.11 ([3]). *Let $(M, \Gamma, U(\Gamma), \xi)$ be a sutured contact 3-manifold with an adapted contact form α , (M^*, α^*) be its completion and J be an almost complex structure on $\mathbb{R} \times M^*$ which is tailored to (M^*, α^*) . Then the contact homology algebra $HC(M, \Gamma, \xi)$ is defined and independent of the choice of contact 1-form α with $\ker(\alpha) = \xi$, adapted almost complex structure J , and abstract perturbation.*

Remark 2.12. Fact 2.6, Theorems 2.7 and 2.8, Formulas 2.2.1 and 2.2.2 hold for J -holomorphic curves in the symplectization of the completion of a sutured contact manifold, provided that we choose the almost complex structure J on $\mathbb{R} \times M^*$ to be tailored to (M^*, α^*) .

Remark 2.13. Observe that Theorem 2.11 and Remark 2.12 rely on the assumption that the machinery, needed to prove the analogous properties for contact homology and cylindrical contact homology in the closed case, works.

3. CONSTRUCTION

3.1. Gluing map. First we construct $H \in C^\infty(\mathbb{R}^2)$. The time-1 flow of the Hamiltonian vector field associated to H composed with an appropriate rotation will play a role of the gluing map when we will apply the gluing construction described in Section 2.1 to the sutured contact solid cylinder constructed in Section 3.2.

We fix $p \in \mathbb{R}^2$ and consider $H_{\text{sing}} : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $H_{\text{sing}} = \mu r^2 \cos(n|k|\theta)$ in polar coordinates about p , where $\mu > 0$, $n \geq 1$ and $k \in \mathbb{Z} \setminus \{-1, 0, 1\}$. Note that H_{sing} is singular only at p . We obtain $H \in C^\infty(\mathbb{R}^2)$ from H_{sing} by perturbing H_{sing} on a disk $D(r_{\text{sing}})$ about p in such a way that H has $n|k|$ equally spaced saddle points, critical point at p and interpolates with no critical points with H_{sing} on $D(r_{\text{sing}})$. In other words, $H = H_{\text{sing}}$ on $\mathbb{R}^2 \setminus D(r_{\text{sing}})$. For the level sets of H_{sing} and H in the case $n = 1$, $|k| = 3$ we refer to Figure 1.

The construction of H is a modification of the construction described in [5].

We proceed in four steps.




FIGURE 1. The level sets of H_{sing} (left) and the level sets of H (right) in the case $n = 1, |k| = 3$

(1) We consider

$$\begin{aligned} H_1 &= H_{sing} + f(r, \theta) = H_{sing} + f_{\exp}(r, \theta) + g(r, \theta) \\ &= \mu r^2 \cos(n|k|\theta) - Ae^{-mr^2} + g(r, \theta), \end{aligned}$$

where A and m are positive constants, and $g(r, \theta)$ is a smooth function to be chosen later. We are interested in the critical points of H_1 away from the origin.

We calculate

$$\begin{aligned} \frac{\partial H_1}{\partial r} &= 2\mu r \cos(n|k|\theta) + 2mrAe^{-mr^2} + \frac{\partial g}{\partial r}, \\ \frac{\partial H_1}{\partial \theta} &= -n|k|\mu r^2 \sin(n|k|\theta). \end{aligned}$$

Thus, at the critical points of H_1 we must have $\sin(n|k|\theta) = 0$. In this case, $\cos(n|k|\theta) = \pm 1$. If $\cos(n|k|\theta) = 1$, then $\frac{\partial H_1}{\partial r} - \frac{\partial g}{\partial r}$ cannot be zero. When $\cos(n|k|\theta) = -1$, $\frac{\partial H_1}{\partial r} - \frac{\partial g}{\partial r} = -2\mu r + 2mrAe^{-mr^2}$. For $r > 0$, $\frac{\partial H_1}{\partial r} - \frac{\partial g}{\partial r} = 0$ when $e^{mr^2} = \frac{mA}{\mu}$, i.e., when $r = r_c := \sqrt{\frac{1}{m} \ln(\frac{mA}{\mu})}$. We impose the restriction that $mA > \mu$. Note that by making m large, we can make r_c arbitrarily small. When $\cos(n|k|\theta) = -1$, $H_1 - g(r, \theta) = -\frac{\mu}{m}(\ln(\frac{mA}{\mu}) + 1)$. Let $g(r)$ be equal to $\frac{\mu}{m}(\ln(\frac{mA}{\mu}) + 1)$ on the annular neighborhood of $r = r_c$. For such g , H_1 is 0 at the critical points, i.e., at the points (r_c, θ) , where $\cos(n|k|\theta) = -1$.

In summary, we get critical points at one value of r at the values of θ when $\cos(n|k|\theta) = -1$, that is, for $n|k|$ values of θ . These are our $n|k|$ saddle points (it's not hard to see they are saddle points; alternatively, we can deduce that they must be for index reasons).

- (2) Keeping f_{exp} solely a function of r and keeping g constant, we cut off f_{exp} smoothly starting at some point past r_c to give a Hamiltonian H_2 which agrees with $H_{sing} + g$ outside a ball. As long as $\frac{\partial f_{exp}}{\partial r} < 2\mu r$, there are no new critical points.

Note that $f_{exp}(r_c) = -\frac{\mu}{m}$. Keeping $\frac{\partial f_{exp}}{\partial r}$ near μr_c (which, using e.g. $A = \frac{e\mu}{m}$, is $\frac{1}{\sqrt{m}}$), we can bring f_{exp} to zero in a radial distance of a constant times $\frac{1}{\sqrt{m}}$; i.e. for m large we can make H_2 agree with $H_{sing} + g$ outside an arbitrarily small ball.

For $A = \frac{e\mu}{m}$, $g = 2\frac{\mu}{m}$. Then keeping g solely a function of r , we cut off $g(r, \theta)$ smoothly starting at some point past the point where $H_2 = H_{sing} + g$ to give Hamiltonian H_3 . As long as $\frac{\partial g}{\partial r} > -2\mu r$, there are no new critical points. We can make it in such a way that H_3 agrees with H_{sing} outside a small ball.

- (3) Recall that $H_3 = H_{sing} + f_{exp} + g$ near the origin and $g(r, \theta) = 2\frac{\mu}{m} > 0$. Note that $g(r, \theta)$ is small for large m . Now keeping g constant we modify $H_{sing} + f_{exp} + g$ near the origin to give us H_4 which is $Br^2 - C$ near the origin (for $B > 0$), which corresponds to the Hamiltonian flow rotating at a constant angular rate. Since $\frac{\partial H_3}{\partial r} = \frac{\partial(H_{sing} + f_{exp})}{\partial r} > 0$ for $r < r_c$, we can patch together $Br^2 - C$ near the origin with H_2 outside a small ball of radius less than r_c in a radially symmetric manner to get H_4 such that $\frac{\partial H_4}{\partial r} > 0$ for $r < r_c$ (we do this by choosing C sufficiently large). Note that H_4 has a critical point at the origin.
- (4) Finally, to ensure no fixed points of the time-1 flow of the Hamiltonian vector field of H , we let H be H_4 multiplied by a radially symmetric function which is ϵ for $r < R$ (for ϵ sufficiently small that the only fixed points of the time-1 flow inside radius R are the critical points and for R large enough that H_4 agrees with H_{sing} for $r > R$) and 1 for $r > 2R$. This creates no new fixed points in the region $R < r < 2R$ because H_4 and $\frac{\partial H_4}{\partial r}$ have the same sign there. Now there are no fixed points of the time-1 flow of the Hamiltonian vector field of H , except for the $n|k|+1$ critical points of H because outside radius R there are no compact flow lines.

Let $p_1, \dots, p_{n|k|}$ denote the equally spaced saddle points of H ordered counterclockwise, i.e., $R_{n|k|}(p_i) = p_{i+1}$, where $R_{n|k|}$ corresponds to the $\frac{2\pi}{n|k|}$ -rotation around the center of $D(r_{sing})$.

Remark 3.1. We first note that $H(p_s) = 0$ for $s = 1, \dots, n|k|$. Hence, by Morse lemma (arguing the same way as in Lemma 3.2 in [8]) we get that there is a neighborhood U_s of p_s such that $H = axy$ on U_s , where $s = 1, \dots, n|k|$ and $a > 0$. In addition, observe that H is $\frac{2\pi}{n|k|}$ -symmetric with respect to θ . Therefore, U_s 's together with coordinates (x, y) are $\frac{2\pi}{n|k|}$ -symmetric with respect to θ , i.e., $R_{n|k|}(U_s) = U_{s+1}$ and coordinates on U_s maps to the coordinate on U_{s+1} , where $R_{n|k|}$ denotes $\frac{2\pi}{n|k|}$ -rotation with respect to θ . Finally, note that $H = \tilde{B}r^2 - \tilde{C}$ on a neighborhood of the center of $D(r_{sing})$, which we call U , where $\tilde{C} > 0$ and $\tilde{B}$ is a small positive number and hence Hamiltonian flow rotates at a constant rate near the origin.

3.2. Sutured contact solid tori. In this section, we construct the sutured contact solid torus with $2n$ sutures of slope $\frac{k}{l}$, where $n \in \mathbb{N}$, $(k, l) = 1$, $|k| > l > 0$.

Let γ_{p, p_s} be an embedded curve in $\mathbb{R}^2$ which starts at p and ends at p_s for $s = 1, \dots, n|k|$. For the time being, we can think about γ_{p, p_s} as about the segment connecting p and p_s .

Lemma 3.2. *There exists a 1-form β on $\mathbb{R}^2$ satisfying the following:*

- (1) $d\beta > 0$;
- (2) its singular foliation given by $\ker \beta$ has isolated singularities and no closed orbits;
- (3) $\beta = \frac{\varepsilon_c}{2}r^2d\theta$ on U with respect to the polar coordinates whose origin is at the center of $D(r_{sing})$; $\beta = \frac{\varepsilon_{sym}}{2}(xdy - ydx)$ on U_s with respect to the coordinates from Remark 3.1, where $s \in \{1, \dots, n|k|\}$; $\beta = \frac{1}{2}r^2d\theta$ on $\mathbb{R}^2 \setminus D(r_{sing})$ with respect to the polar coordinates whose origin is at the center of $D(r_{sing})$; here $0 < \varepsilon_c \ll \varepsilon_{sym} \ll 1$;
- (4) the set of hyperbolic points of the singular foliation of β is given by $\{q_s\}_{s=1}^{n|k|}$ such that q_s lies on γ_{p,p_s} outside of U_s and U ;
- (5) β is $\frac{2\pi}{n|k|}$ -symmetric, i.e., $R_{n|k|}^*(\beta) = \beta$, where $R_{n|k|} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a $\frac{2\pi}{n|k|}$ -rotation with respect to the center of $D(r_{sing})$.

Proof. Consider a singular foliation $\mathcal{F}$ on $\mathbb{R}^2$ which satisfies the following:

- (1) $\mathcal{F}$ is Morse-Smale and has no closed orbits.
- (2) The singular set of $\mathcal{F}$ consists of elliptic points and hyperbolic points. The elliptic points are the equally spaced saddle points of H and the center of $D(r_{sing})$. The set of hyperbolic points of the singular foliation of β is given by $\{q_s\}_{s=1}^{n|k|}$ such that q_s lies on γ_{p,p_s} outside of U_s and U .
- (3) $\mathcal{F}$ is oriented and for one choice of orientation the flow is transverse to and exits from $\partial D(r_{sing})$.
- (4) $\mathcal{F}$ is $\frac{2\pi}{n|k|}$ -symmetric with respect to θ .

Next, we modify $\mathcal{F}$ near each of the singular points so that $\mathcal{F}$ is given by $\beta_0 = \frac{1}{2}(xdy - ydx)$ on U_s with respect to the coordinates from Remark 3.1 and $\beta_0 = 2xdy + ydx$ near a hyperbolic point. On $\mathbb{R}^2 \setminus D(r_{sing})$, $\beta_0 = \frac{1}{2}r^2d\theta$ with respect to the polar coordinates whose origin is at the center of $D(r_{sing})$. In addition, on U , $\beta_0 = \frac{1}{2}r^2d\theta$ with respect to the polar coordinates whose origin is at the center of $D(r_{sing})$. From Remark 3.1 it follows that we can do it in such a way that the modification of $\mathcal{F}$ is still $\frac{2\pi}{n|k|}$ -symmetric. Finally, we get $\mathcal{F}$ given by β_0 , which satisfies $d\beta_0 > 0$ near the singular points and on $\mathbb{R}^2 \setminus D(r_{sing})$. Now let $\beta = g\beta_0$, where g is a positive function with $dg(X) \gg 0$ outside of $U \cup (\cup_{s=1}^{n|k|} U_s) \cup (\mathbb{R}^2 \setminus D(r_{sing}))$, $g|_{\cup_{s=1}^{n|k|} U_s} = \varepsilon_{sym}$, $g|_U = \varepsilon_c$, $g|_{\mathbb{R}^2 \setminus D(r_{sing})} = 1$ and X is an oriented vector field for $\mathcal{F}$ (nonzero away from the singular points). Here $0 < \varepsilon_c \ll \varepsilon_{sym} \ll 1$. Since $d\beta = dg \wedge \beta_0 + g \wedge d\beta_0$, $dg(X) \gg 0$ guarantees that $d\beta > 0$. $\square$

Remark 3.3. From the previous lemma we get β defined on $\mathbb{R}^2$ with the following properties:

- (i) $d\beta > 0$ on $\mathbb{R}^2$;
- (ii) $\beta = \frac{\varepsilon_c}{2}r^2d\theta$ and $H = \tilde{B}r^2 - \tilde{C}$ on U , where $\tilde{C} > 0$; here ε_c is a small positive real number;
- (iii) $\beta = \frac{\varepsilon_{sym}}{2}(xdy - ydx)$ and $H = axy$ on U_s for $s = 1, \dots, n|k|$; here $0 < \varepsilon_c \ll \varepsilon_{sym} \ll 1$;
- (iv) $\beta = \frac{1}{2}r^2d\theta$ and $H = \mu r^2 \cos(n\theta)$ on $\mathbb{R}^2 \setminus D(r_{sing})$.

For the comparison of the level sets of H with the singular foliation of β in the case $n = 1$, $|k| = 3$ we refer to Figure 2.

Lemma 3.4. Let β be a 1-form from Lemma 3.2. The Hamiltonian vector field X_H of H with respect to the area form $d\beta$ satisfies $\beta(X_H) = H$ on $(\cup_{s=1}^{n|k|} U_s) \cup (\mathbb{R}^2 \setminus D(r_{sing}))$. In addition, the Hamiltonian vector field X_H of H with respect to the area form $d\beta$ satisfies $\beta(X_H) - H = \tilde{C}$ on U .




FIGURE 2. The level sets of H (left) and the characteristic foliation of β (right) in the case $n = 1, |k| = 3$

Proof. First, Remark 3.3 implies that $\beta = \frac{\varepsilon_c}{2}r^2d\theta$, $H = \tilde{B}r^2 - \tilde{C}$ on U and ε_c is a small positive number. Now we show that $X_H = \frac{2\tilde{B}}{\varepsilon_c}\frac{\partial}{\partial\theta}$ is a solution of $\beta(X_H) - H = \tilde{C}$ on U . We calculate

$$i_{X_H}(d\beta) = \left(\frac{2\tilde{B}}{\varepsilon_c}\frac{\partial}{\partial\theta}\right) \lrcorner (\varepsilon_c r dr \wedge d\theta) = -2\tilde{B}r dr = -dH,$$

and

$$\beta(X_H) - H = \left(\frac{\varepsilon_c}{2}r^2d\theta\right) \left(\frac{2\tilde{B}}{\varepsilon_c}\frac{\partial}{\partial\theta}\right) - \tilde{B}r^2 + \tilde{C} = \tilde{C}.$$

Next, we work on U_s , where $s = 1, \dots, n|k|$. From Remark 3.3 it follows that $\beta = \frac{\varepsilon_{sym}}{2}(xdy - ydx)$ and $H = axy$ on U_s . Let X_H be a Hamiltonian vector field defined by $i_{X_H}d\beta = -dH$.

We show that

$$X_H = -\frac{ax}{\varepsilon_{sym}}\frac{\partial}{\partial x} + \frac{ay}{\varepsilon_{sym}}\frac{\partial}{\partial y}$$

is a solution of the equation

$$(3.2.1) \quad \beta(X_H) = H$$

on U_s . We calculate

$$i_{X_H}(d\beta) = \left(-\frac{ax}{\varepsilon_{sym}}\frac{\partial}{\partial x} + \frac{ay}{\varepsilon_{sym}}\frac{\partial}{\partial y}\right) \lrcorner (\varepsilon_{sym}dx \wedge dy) = -axdy - aydx = -dH$$

and

$$\beta(X_H) = \frac{\varepsilon_{sym}}{2}(xdy - ydx) \left(-\frac{ax}{\varepsilon_{sym}} \frac{\partial}{\partial x} + \frac{ay}{\varepsilon_{sym}} \frac{\partial}{\partial y} \right) = axy = H.$$

Finally, Remark 3.3 says that $\beta = \frac{1}{2}r^2 d\theta$ and $H = \mu r^2 \cos(n|k|\theta)$ on $\mathbb{R}^2 \setminus D(r_{sing})$. As in the previous case, we show that

$$X_H = n|k|\mu r \sin(n|k|\theta) \frac{\partial}{\partial r} + 2\mu \cos(n|k|\theta) \frac{\partial}{\partial \theta}$$

is a solution of Equation (3.2.1) on $\mathbb{R}^2 \setminus D(r_{sing})$.

We calculate

$$\begin{aligned} i_{X_H}(d\beta) &= (n|k|\mu r \sin(n|k|\theta) \partial_r + 2\mu \cos(n|k|\theta) \partial_\theta) \lrcorner (rdr \wedge d\theta) \\ &= -2\mu r \cos(n|k|\theta) dr + n|k|\mu r^2 \sin(n|k|\theta) d\theta = -dH, \end{aligned}$$

and

$$\begin{aligned} \beta(X_H) &= \left(\frac{1}{2}r^2 d\theta \right) \left(n|k|\mu r \sin(n|k|\theta) \frac{\partial}{\partial r} + 2\mu \cos(n|k|\theta) \frac{\partial}{\partial \theta} \right) \\ &= \mu r^2 \cos(n|k|\theta) = H. \end{aligned}$$

□

Let X_H be the Hamiltonian vector field of H with respect to $d\beta$ and $\varphi_{X_H}^s$ be the time- s flow of X_H . Now we introduce the following notations:

$$\begin{aligned} S &:= \{x \in \mathbb{R}^2 \setminus D(r_{sing}) \mid \varphi_{X_H}^s(x) \in \mathbb{R}^2 \setminus D(r_{sing}) \forall s \in [0, 1]\}, \\ V &:= \{x \in U \mid \varphi_{X_H}^s(x) \in U \forall s \in [-1, 1]\}, \quad \text{and} \\ V_i &:= \{x \in U_i \mid \varphi_{X_H}^s(x) \in U_i \forall s \in [-1, 1]\}. \end{aligned}$$

For simplicity, let us denote $\varphi_{X_H} := \varphi_{X_H}^1$.

Remark 3.5. Using the form of X_H on U_i , where $i = 1, \dots, n|k|$, we may assume that the curves γ_{p,p_i} 's in Lemma 3.2 satisfy the following list of properties:

- (1) γ_{p,p_i} is an embedded curve which starts at p and ends at p_i ;
- (2) γ_{p,p_i} is a part of one of the curves of the singular foliation given by $\ker \beta$;
- (3) γ_{p,p_i} coincides with one of the level sets of H on V_i and near p_i can be presented as $W^s(\varphi_{X_H}, p_i) = \{x \mid (\varphi_{X_H})^n(x) \rightarrow p \text{ as } n \rightarrow \infty\}$.

Recall that the following claim was proven in [8]:

Claim 3.6 ([8]). *If (M, ω) is an exact symplectic manifold, i.e., $\omega = d\beta$, then the flow $\varphi_{X_H}^t$ of a Hamiltonian vector field X_H consists of exact symplectic maps, i.e.,*

$$(\varphi_{X_H}^t)^* \beta - \beta = df_t,$$

where

$$f_t = \int_0^t (-H + \beta(X_H)) \circ \varphi_{X_H}^s ds.$$

Remark 3.7. Observe that from Lemma 3.4 and Claim 3.6 it follows that $\varphi_{X_H}^*(\beta) - \beta = dh$, where $h := f_1 = 0$ on $S \cup (\cup_{i=1}^{n|k|} V_i)$ and $h = \tilde{C} > 0$ on V . Hence, we get $\varphi_{X_H}^*(\beta) = \beta$ on $S \cup V \cup (\cup_{i=1}^{n|k|} V_i)$.

Now we define $\varphi_{\frac{k}{l}} := R_{-\frac{k}{l}} \circ \varphi_{X_H}$, where $R_{-\frac{k}{l}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a $-\frac{2\pi l}{k}$ -rotation around the center of $D(r_{\text{sing}})$.

Remark 3.8. Since $R_{n|k|}^*(\beta) = \beta$, we get $R_{-\frac{k}{l}}^*(\beta) = \beta$ and hence

$$\varphi_{\frac{k}{l}}^*(\beta) = (R_{-\frac{k}{l}} \circ \varphi_{X_H})^*(\beta) = \varphi_{X_H}^*(R_{-\frac{k}{l}}^*(\beta)) = \varphi_{X_H}^*(\beta).$$

Fix $R_* \gg r_{\text{sing}}$ such that there is an annular neighborhood V_{R_*} of $\partial D(R_*)$ in $\mathbb{R}^2$ with $V_{R_*} \subset S$. Consider $D(R_*)$ with $\beta_0 := \beta|_{D(R_*)}$ and $\beta_1 := \varphi_{X_H}^*(\beta)|_{D(R_*)} (= \varphi_{\frac{k}{l}}^*(\beta)|_{D(R_*)})$. Note that

$$(3.2.2) \quad d\beta_1 = d(\varphi_{X_H}^*(\beta)|_{D(R_*)}) = \varphi_{X_H}^*(d\beta)|_{D(R_*)} = (d\beta)|_{D(R_*)} = d\beta_0 > 0.$$

In addition, from the definitions of $V(R_*)$ and $D(R_*)$ it follows that

$$(3.2.3) \quad \beta_0 = \beta_1 \quad \text{on} \quad V_{R_*} \cap D(R_*).$$

Now we recall Lemma 3.10 from [8], which provides the construction of the contact 1-form on $[-1, 1] \times D^2$.

Lemma 3.9 ([8]). *Let β_0 and β_1 be two 1-forms on D^2 such that $\beta_0 = \beta_1$ in a neighborhood of ∂D^2 and $d\beta_0 = d\beta_1 = \omega > 0$. Then there exists a contact 1-form α and a Reeb vector field R_α on $[-1, 1] \times D^2$ with coordinates (t, x) , where t is a coordinate on $[-1, 1]$ and x is a coordinate on D^2 , with the following properties:*

- (1) $\alpha = dt + \varepsilon\beta_0$ in a neighborhood of $\{-1\} \times D^2$;
- (2) $\alpha = dt + \varepsilon\beta_1$ in a neighborhood of $\{1\} \times D^2$;
- (3) R_α is collinear to $\frac{\partial}{\partial t}$ on $[-1, 1] \times D^2$;
- (4) $R_\alpha = \frac{\partial}{\partial t}$ in a neighborhood of $[-1, 1] \times \partial D^2$.

Here ε is a small positive number.

In addition, recall that

$$(3.2.4) \quad \alpha = (1 + \varepsilon\chi_1(t)h)dt + \varepsilon((1 - \chi_0(t))\beta_0 + \chi_0(t)\beta_1),$$


where $h \in C^\infty(D^2)$ such that $\beta_1 - \beta_0 = dh$; $\chi_0 : [-1, 1] \rightarrow [0, 1]$ is a smooth map for which $\chi_0(t) = 0$ for $-1 \leq t \leq -1 + \varepsilon_{\chi_0}$, $\chi_0(t) = 1$ for $1 - \varepsilon_{\chi_0} \leq t \leq 1$, $\chi_0'(t) \geq 0$ for $t \in [-1, 1]$ and ε_{χ_0} is a small positive number; $\chi_1(t) := \chi_0'(t)$; ε is a sufficiently small positive number.

Remark 3.10. Note that $d\alpha = \varepsilon\omega$, where α is a 1-form given by Formula 3.2.4 and $\omega = d\beta_0 = d\beta_1 > 0$ on D^2 .

Observe that from Formulas 3.2.2 and 3.2.3 it follows that β_0 and β_1 described above satisfy the conditions of Lemma 3.9. We now take $[-1, 1] \times D(R_*)$ equipped with the contact 1-form α given by Formula 3.2.4. For simplicity, let us denote $\beta_- := \varepsilon\beta_0$ and $\beta_+ := \varepsilon\beta_1$, where ε is a constant from Lemma 3.9 which makes α contact.

Then we construct P_+ , P_- and D in the way described in [8]. Recall that

$$P_+, P_-, D \subset D(R_*) \subset \mathbb{R}^2$$

FIGURE 3. Construction of P_+ , P_- and D in the case $n = 1$, $|k| = 3$

are surfaces with boundary which satisfy the following properties:

- (1) $P_{\pm} \subset D$;
- (2) $(\partial P_{\pm})_{\partial} \subset \partial D$ and $(\partial P_{\pm})_{int} \subset int(D)$;
- (3) φ_{X_H} maps P_+ to P_- in such a way that $\varphi_{X_H}((\partial P_+)_{int}) = (\partial P_-)_{\partial}$ and $\varphi_{X_H}((\partial P_+)_{\partial}) = (\partial P_-)_{int}$;
- (4) $(\partial P_-)_{\partial} \cap (\partial P_+)_{\partial} = \emptyset$.

Note that in our case

- $\partial P_+ = (\cup_{s=0}^{n|k|-1} a_s^+) \cup (\cup_{s=0}^{n|k|-1} b_s^+)$,
- $\partial P_- = (\cup_{s=0}^{n|k|-1} a_s^-) \cup (\cup_{s=0}^{n|k|-1} b_s^-)$,
- $\partial D = (\cup_{s=0}^{n|k|-1} a_s^+) \cup (\cup_{s=0}^{n|k|-1} b_s^-) \cup (\cup_{s=0}^{n|k|-1} c_s^+) \cup (\cup_{s=0}^{n|k|-1} c_s^-)$,

which is slightly different from the case described in [8] when

- $\partial P_+ = (\cup_{s=0}^{n-1} a_s^+) \cup (\cup_{s=0}^{n-1} b_s^+)$,
- $\partial P_- = (\cup_{s=0}^{n-1} a_s^-) \cup (\cup_{s=0}^{n-1} b_s^-)$,
- $\partial D = (\cup_{s=0}^{n-1} a_s^+) \cup (\cup_{s=0}^{n-1} b_s^-) \cup (\cup_{s=0}^{n-1} c_s^+) \cup (\cup_{s=0}^{n-1} c_s^-)$.

See Figure 3 for the schematic visualization of P_+ (bounded by the bold line), P_- and D . For more details of this construction we refer to [8].

Remark 3.11. Note that $a_i^{\pm}$'s, $b_i^{\pm}$'s and $c_i^{\pm}$'s are constructed in such a way that

$$a_i^{\pm}, b_i^{\pm}, c_i^{\pm} \subset D(R_*) \cap S$$

for $i = 0, \dots, n|k| - 1$. Hence, we see that $\partial P_+, \partial P_-, \partial D \subset D(R_*) \cap S$. In addition, $R_{n|k|}(a_i^{\pm}) = a_{i+1}^{\pm}$ and $R_{n|k|}(b_i^{\pm}) = b_{i+1}^{\pm}$, where $R_{n|k|}$ is a $\frac{2\pi}{n|k|}$ -rotation around p and $i, i+1$ are considered modulo $n|k|$.

We take $[-1, 1] \times D$ with a contact form $\alpha := \alpha|_{[-1, 1] \times D}$. Let $\Gamma = \{0\} \times \partial D$ in $[-1, 1] \times D$ and $U(\Gamma) := [0, 1] \times [-1, 1] \times \Gamma$ be a neighborhood of Γ with coordinates $(\tau, t) \in [0, 1] \times [-1, 1]$,

where t is a usual t -coordinate on $[-1, 1] \times D$. From the definition of S and Remark 3.11 it follows that we may assume that $U(\Gamma) \subset [-1, 1] \times (S \cap D)$.

Lemma 3.12. *$([-1, 1] \times D, \Gamma, U(\Gamma), \xi)$ is a sutured contact manifold and α is an adapted contact form.*

Proof. First note that $\alpha|_{R_-} = \beta_-$ and $\alpha|_{R_+} = \beta_+$. Let us check that (R_-, β_-) and (R_+, β_+) are Liouville manifolds. From the construction of $\beta_{\pm}$ it follows that $d(\beta_-) = d(\beta_+) > 0$. Since $\beta_- = \beta_+$ on $D \cap S$ and by Formula 3.2.4, $\alpha = dt + \beta_-$ on $U(\Gamma)$. Recall that $\beta_- = \beta_+ = \frac{\varepsilon}{2}r^2d\theta$ on $D \cap S$. Hence, $\alpha|_{U(\Gamma)} = dt + \frac{\varepsilon}{2}r^2d\theta$. The calculation

$$i_{Y_{\pm}|_{R_{\pm} \cap U(\Gamma)}}(d\beta_{\pm}) = \left(\frac{1}{2}r\partial_r\right) \lrcorner (\varepsilon r dr \wedge d\theta) = \frac{\varepsilon}{2}r^2d\theta = \beta_{\pm}$$

implies that the Liouville vector fields $Y_{\pm}|_{R_{\pm} \cap U(\Gamma)}$ are equal to $\frac{1}{2}r\partial_r$. From the construction of D it follows that $Y_{\pm}$ is positively transverse to $\partial R_{\pm}$. Therefore, $(R_-, \varepsilon\beta_0)$ and $(R_+, \varepsilon\beta_1)$ are Liouville manifolds. As we already mentioned, $\alpha = dt + \beta_-$ on $U(\Gamma)$. Finally, if we take τ such that $\partial_{\tau} = \frac{1}{2}r\partial_r$, then $([-1, 1] \times D, \Gamma, U(\Gamma), \xi)$ becomes a sutured contact manifold with an adapted contact form α . $\square$

Then we use $\varphi_{\frac{k}{l}}$ for the gluing construction. Note that φ_{X_H} maps a_s^+ to a_s^- and b_s^+ to b_s^- . Hence, using Remark 3.11, we see that $\varphi_{\frac{k}{l}}$ maps a_s^+ to a_{s-nl}^- and b_s^+ to b_{s-nl}^- for $k > 0$, and $\varphi_{\frac{k}{l}}$ maps a_s^+ to a_{s+nl}^- and b_s^+ to b_{s+nl}^- for $k < 0$. Then we follow the gluing procedure briefly described in Section 2.1 and completely written in [3]. Finally, we get a sutured contact solid torus $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}))$ with a contact form $\tilde{\alpha}_{\delta}$, where $\tilde{\Gamma}$ is a set of $2n$ parallel closed curves of slope $\frac{k}{l}$, where $n \in \mathbb{N}$, $(k, l) = 1$, $|k| > l > 0$ and δ is the rotation angle of the map φ_{X_H} near p .

Remark 3.13. We have constructed $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}))$ using the gluing construction for sutured manifolds. However, since there is a close connection between sutured contact manifolds and contact manifolds with convex boundary, we observe that the gluing construction we used for the sutured contact solid cylinder corresponds to the gluing construction for the contact 3-ball with convex boundary and one dividing curve on the boundary. The corresponding gluing construction for the contact 3-ball with convex boundary corresponds (is inverse) to the convex decomposition of the contact solid torus $S^1 \times D^2$ with convex boundary with respect to the convex meridional disk $\{pt\} \times D^2$ with ∂ -parallel dividing curves. Hence, the constructed sutured contact solid tori are universally tight sutured contact manifolds by the gluing/classification result from Section 2 in [9] (more precisely, Corollary 2.3, Theorem 2.5 and Corollary 2.6).

4. CALCULATION

4.1. Reeb orbits. Note that $\varphi_{\frac{k}{l}}|_{P_+}$ has n orbits of period $|k|$ obtained from the equally spaced saddle points of H . Lemma 3.9 and the gluing procedure briefly described in Section 2.1 imply that these orbits correspond to the Reeb orbits, which we call $\gamma_1, \dots, \gamma_n$ such that

$$[\gamma_s] = [\gamma_t] = |k| \in H_1(S^1 \times D^2; \mathbb{Z})$$

for $s, t = 1, \dots, n$. In addition, $\varphi_{\frac{k}{l}}|_{P_+}$ has a periodic point of period 1, which is p . It corresponds to the Reeb orbit, which we call γ , such that $[\gamma] = 1 \in H_1(S^1 \times D^2; \mathbb{Z})$.

Lemma 4.1. $\int_{\gamma_s} \tilde{\alpha}_\delta = \int_{\gamma_t} \tilde{\alpha}_\delta$ and $|k| \int_{\gamma} \tilde{\alpha}_\delta > \int_{\gamma_s} \tilde{\alpha}_\delta$, where $s, t = 1, \dots, n$.

Proof. Let

$$M^{(0)} = (([-1, 1] \times D) \cup (R_+(\Gamma) \times [1; \infty)) \cup (R_+(\Gamma) \times (-\infty; -1]))$$

and

$$\tilde{M} = M^{(0)} \setminus ((P_+ \times (N, \infty) \cup (P_- \times (-\infty, -N))).$$

In addition, let $\alpha_{\tilde{M}}$ denote the contact form on $\tilde{M}$ and let $\xi_{\tilde{M}}$ denote the contact structure defined by $\alpha_{\tilde{M}}$.

Consider $[-1, 1] \times D \subset \tilde{M}$. From the construction of α it follows that $\beta_+ = \beta_-$ on V_s and $\alpha|_{[-1, 1] \times V_s} = dt + \beta_-$ for $s = 1, \dots, n|k|$. Hence, since the contact structure on $[1, \infty) \times P_+$ is given by $dt + \beta_+$ and the contact structure on $(-\infty, -1] \times P_-$ is given by $dt + \beta_-$, $\alpha_{\tilde{M}}|_{[-N, N] \times V_s} = dt + \beta_-$ on $[-N, N] \times V_s \subset \tilde{M}$ for $s = 1, \dots, n|k|$. Therefore, we get

$$(4.1.1) \quad \int_{[-N, N] \times \{p_s\}} \alpha_{\tilde{M}} = 2N$$

for $s = 1, \dots, n|k|$. From the gluing construction and Equation (4.1.1) it follows that

$$\int_{\gamma_s} \tilde{\alpha}_\delta = 2N|k|$$

for $s = 1, \dots, n$. Note that $\int_{\gamma_s} \tilde{\alpha}_\delta$ does not depend on s . Hence, $\int_{\gamma_s} \tilde{\alpha}_\delta = \int_{\gamma_t} \tilde{\alpha}_\delta$ for $s, t = 1, \dots, n$.

Now from the fact that $\alpha = (1 + \varepsilon \chi_1(t)h)dt + \beta_-$ on $[-1, 1] \times V$, where $h > 0$ and $\chi_1(t) > 0$, we get that

$$R_\alpha = \frac{1}{1 + \varepsilon \chi_1(t)h} \frac{\partial}{\partial t}$$

on $[-1, 1] \times V$. Hence, from the gluing construction we obtain $|k| \int_{\gamma} \tilde{\alpha}_\delta > 2N|k|$. Thus,

$$\int_{\gamma_s} \tilde{\alpha}_\delta = \int_{\gamma_t} \tilde{\alpha}_\delta \quad \text{and} \quad |k| \int_{\gamma} \tilde{\alpha}_\delta > \int_{\gamma_s} \tilde{\alpha}_\delta,$$

where $s, t = 1, \dots, n$. □

For simplicity, from now on we assume that $l = 1$. The calculation for the case when $l > 1$ is completely analogous.

Lemma 4.2. *All closed orbits of $R_{\tilde{\alpha}_\delta}$ are nondegenerate. Moreover, γ is an elliptic orbit and γ_i is a hyperbolic orbit such that γ^t and γ_i^s are good orbits for $i = 1, \dots, n$; $s, t \in \mathbb{N}$. There exists a symplectic trivialization τ of ξ along γ and γ_i 's constructed in the consistent way as described in Section 2.2, and $N_\delta \in \mathbb{N}$ such that*

$$\mu_\tau(\gamma_i^s) = \begin{cases} 2s, & \text{for } k < -1; \\ -2s, & \text{for } k > 1, \end{cases}$$

and

$$\mu_\tau(\gamma^t) = \begin{cases} 2m - 1, & \text{for } k < -1 \text{ and } (m-1)|k| \leq t < m|k|; \\ -2m + 1, & \text{for } k > 1 \text{ and } (m-1)|k| < t \leq m|k| \end{cases}$$

for $i = 1, \dots, n$; $t \leq N_\delta$, $s \leq \frac{N_\delta}{|k|}$.

Proof. Fix $i = 1, \dots, n$. We first observe that $H|_{V_i} = axy$, where $a > 0$ and hence

$$\varphi_{X_H}|_{V_i} = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix},$$

where $\lambda = e^a \neq 1$. Let the symplectic trivialization of $\xi_{\tilde{M}}$ along $[-N, N] \times \{p_i\}$ be given by the framing $(\lambda^{\frac{-N-t}{2N}} \partial_x, \lambda^{\frac{t+N}{2N}} \partial_y)$, where $i = 1, \dots, n|k|$ and (x, y) are coordinates on V_i which coincide with the coordinates on U_i from Remark 3.1. Since Remark 3.1 implies that $R_{n|k|}$ maps coordinates on V_i to the coordinate on V_{i+1} , where $i, i+1$ are considered modulo $n|k|$ and $R_{n|k|}$ denotes $\frac{2\pi}{n|k|}$ -rotation around p , we conclude that the symplectic trivializations of $\xi_{\tilde{M}}$ along $[-N, N] \times \{p_{i+nm}\}$'s for $m = 0, \dots, |k| - 1$ and fixed $i = 1, \dots, n$ give rise to the symplectic trivialization τ_{γ_i} of ξ along γ_i . It is easy to see that the linearized return map P_{γ_i} with respect to this trivialization is given by

$$P_{\gamma_i} = \begin{pmatrix} \lambda^{|k|} & 0 \\ 0 & \lambda^{-|k|} \end{pmatrix}.$$

Since the eigenvalues of P_{γ_i} are positive real numbers different from 1, γ_i is a positive hyperbolic orbit. In addition, $P_{\gamma_i^s} = P_{\gamma_i}^s$. Therefore, the eigenvalues of $P_{\gamma_i^s}$ are different from 1. Hence, γ_i^s is a nondegenerate orbit for $s \in \mathbb{N}$ and $i = 1, \dots, n$. We now observe that the linearized Reeb flow around γ_i (with respect to τ_{γ_i}) rotates the eigenspaces of P_{γ_i} by angle 2π for $k < -1$ and -2π for $k > 1$. Hence, we get

$$(4.1.2) \quad \mu_{\tau_{\gamma_i}}(\gamma_i^s) = \begin{cases} 2s, & \text{for } k < -1; \\ -2s, & \text{for } k > 1 \end{cases}$$

for $s \in \mathbb{N}$ and $i = 1, \dots, n$.

Now let the symplectic trivialization of $\xi_{\tilde{M}}$ along $[-N, N] \times \{p\}$ be given by the framing

$$(\cos(\theta_{\delta,k,N}(t))\partial_x + \sin(\theta_{\delta,k,N}(t))\partial_y, -\sin(\theta_{\delta,k,N}(t))\partial_x + \cos(\theta_{\delta,k,N}(t))\partial_y),$$

where $\theta_{\delta,k,N}(t) = \frac{\pi(1-\delta k)(t+N)}{Nk}$ and $t \in [-N, N]$. Observe that $R_{-k} \circ \varphi_{X_H}|_V$ is a rotation by angle $2\pi(-\frac{1}{k} + \delta)$, where R_{-k} is a $-\frac{2\pi}{k}$ -rotation about p and δ is a small positive irrational number. It is easy to see that with respect to this framing P_γ is a rotation by $2\pi(-\frac{1}{k} + \delta)$. Hence, since δ is irrational, we see that γ is an elliptic orbit and γ^t is nondegenerate for $t \in \mathbb{N}$. Let

$$N_\delta := \max\{m \in \mathbb{N} \mid m\delta < \frac{1}{|k|}\}.$$

Note that we get

$$(4.1.3) \quad \mu_{\tau_\gamma}(\gamma^t) = \begin{cases} 2m - 1, & \text{for } k < -1 \text{ and } (m-1)|k| \leq t < m|k|; \\ -2m + 1, & \text{for } k > 1 \text{ and } (m-1)|k| < t \leq m|k| \end{cases}$$

for $t \leq N_\delta$. Formulas 4.1.2 and 4.1.3 and the fact that δ is irrational imply that the parity of $\mu_{\tau_{\gamma_i}}(\gamma_i^s)$ is independent of s for given i and the parity of $\mu_{\tau_\gamma}(\gamma^t)$ is independent of t . Hence, we conclude that γ_i^s 's and γ^t 's are good Reeb orbits for $i = 1, \dots, n$ and $s, t \in \mathbb{N}$.

We now show that the symplectic trivializations τ_{γ_i} 's and τ_γ are consistent in the sense of Section 2.2. We first consider $\{-N\} \times P_- \subset \tilde{M}$. Note that Remark 3.5 implies that for $i = 1, \dots, n|k|$ there is an embedded curve γ_{p,p_i} with the following properties:

- (1) γ_{p,p_i} is an embedded curve which starts at p and ends at p_i ;
- (2) γ_{p,p_i} coincides with one of the level sets of H on V_i and near p_i can be presented as $W^s(\varphi_{X_H}, p_i) = \{x \mid (\varphi_{X_H})^n(x) \rightarrow p \text{ as } n \rightarrow \infty\}$;
- (3) γ_{p,p_i} is a part of one of the curves of the characteristic foliation of $\alpha_{\tilde{M}}$ on $\{-N\} \times P_- \subset \tilde{M}$.

Now consider γ_{p,p_i} with a vector field $v_i \in T\gamma_{p,p_i}$ such that v_i does not vanish and $(R_{n|k|})_*(v_i) = v_{i+1}$, where $R_{n|k|}$ denotes $\frac{2\pi}{n|k|}$ -rotation around p and $i, i+1$ are considered modulo $n|k|$. Let

$$v_{i,t} := -\varepsilon \frac{\beta_t \left(\left(\varphi_{X_H}^{\frac{-N-t}{2N}} \right)_* (v_i) \right)}{1 + \varepsilon h \chi_{1,ext}(t)} \partial_t + \left(\varphi_{X_H}^{\frac{-N-t}{2N}} \right)_* (v_i),$$

where $\beta_t = (1 - \chi_{0,ext}(t))\beta_0 + \chi_{0,ext}(t)\beta_1$,

$$\chi_{0,ext}(t) := \begin{cases} 1, & \text{for } t \in [1, N]; \\ \chi_0(t), & \text{for } t \in [-1, 1]; \\ 0, & \text{for } t \in [-N, -1], \end{cases}$$

and $\chi_{1,ext}(t) := \chi'_{0,ext}(t)$ on $[-N, N]$. It is easy to see that

$$\begin{aligned} \alpha_{\tilde{M}}(v_{i,t}) &= ((1 + \varepsilon \chi_{1,ext}(t)h)dt + \varepsilon \beta_t) \left(-\varepsilon \frac{\beta_t \left(\left(\varphi_{X_H}^{\frac{-N-t}{2N}} \right)_* (v_i) \right)}{1 + \varepsilon h \chi_{1,ext}(t)} \partial_t + \left(\varphi_{X_H}^{\frac{-N-t}{2N}} \right)_* (v_i) \right) \\ &= 0 \end{aligned}$$

along $\varphi^{(-N-t)/2N}(\gamma_{p,p_i})$, where $t \in [-N, N]$ and $i = 1, \dots, n|k|$. Hence, we conclude that $v_{i,t} \in \xi_{\tilde{M}}$ along $\varphi^{(-N-t)/2N}(\gamma_{p,p_i})$ for $i = 1, \dots, n|k|$; $t \in [-N, N]$. In addition, observe that $(\varphi_{X_H}^{(-N-t)/2N})_*(v_i) \in T\varphi^{(-N-t)/2N}(\gamma_{p,p_i})$ does not vanish, since v_i does not vanish. Thus, $v_{i,t}$ does not vanish as well. We now note that Properties (2) and (3) imply that $((\varphi_{X_H}^{(-N-t)/2N})_* v_i)|_{V \cup V_i} \in \xi_{\tilde{M}}$. Hence, we conclude that $\beta_t((\varphi_{X_H}^{(-N-t)/2N})_*(v_i)|_{V \cup V_i}) = 0$ for $t \in [-N, N]$. Thus,

$$v_{i,t} = (\varphi_{X_H}^{(-N-t)/2N})_*(v_i)$$

on $[-N, N] \times (V \cup V_i)$. In addition, Property (2), Property (3) and the fact that p and p_i are fixed points of $\varphi_{X_H}^{(-N-t)/2N}$ for $t \in [-N, N]$ imply that

$$(4.1.4) \quad ((\varphi_{X_H}^{(-N-t)/2N})_*(v_i))(p_i) = h(t)v(p_i),$$

where $h(t)$ does not vanish for $t \in [-N, N]$. Besides that, observe that Property (2) implies that $v(p_i)$ is a multiple of a coordinate vector on V_i . Then we take $(v_{i,t}, w_{i,t})$ as a symplectic basis of $\xi_{\tilde{M}}$ along $(\varphi_{X_H}^{(-N-t)/2N})(\gamma_{p,p_i})$, where $w_{i,t}$ is a symplectic complement of $v_{i,t}$ with respect to $d\alpha_{\tilde{M}}$. We now observe that Remark 3.10 and the fact that $\alpha_{\tilde{M}}|_{[1,N] \times (V \cup V_i)} = dt + \beta_+$, $\alpha_{\tilde{M}}|_{[-N,-1] \times (V \cup V_i)} =$

$dt + \beta_-$ imply that $d\alpha_{\tilde{M}}|_{[-N, N] \times (V \cup V_i)} = \varepsilon\omega$. Hence, $d\alpha_{\tilde{M}}$ is invariant under $(\varphi_{X_H}^{(t+N)/2N})^*$ in the neighborhoods of $[-N, N] \times \{p_i\}$ and $[-N, N] \times \{p\}$ for $i = 1, \dots, n|k|$, $t \in [-N, N]$. Therefore, using the fact that $v_{i,t} = (\varphi_{X_H}^{(-N-t)/2N})_*(v_i)$ on $[-N, N] \times (V \cup V_i)$, we get that $w_{i,t} = (\varphi_{X_H}^{(-N-t)/2N})_*(w_i)$ on $[-N, N] \times (V \cup V_i)$, where $w_i := w_{i,t}|_{\gamma_{p,p_i}}$ and $t \in [-N, N]$. Now we take the symplectic trivialization given by the framing

$$(4.1.5) \quad (v_{i,t}(p_i), w_{i,t}(p_i)) = ((\varphi_{X_H}^{(-N-t)/2N})_*(v_i)(p_i), (\varphi_{X_H}^{(-N-t)/2N})_*(w_i)(p_i))$$

along $[-N, N] \times \{p_i\}$, where $i = 1, \dots, n|k|$. In addition, for a fixed $i = 1, \dots, n$ we consider the sequence of framings

$$(v_{i+nm,t}(p), w_{i+nm,t}(p))_{m=0}^{|k|-1}$$

along $[-N, N] \times \{p\}$. We can rewrite it as

$$(4.1.6) \quad (((\varphi_{X_H}^{(-N-t)/2N})_*(v_{i+nm})(p), (\varphi_{X_H}^{(-N-t)/2N})_*(w_{i+nm})(p)))_{m=0}^{|k|-1}.$$

Finally, note that since

$$(R_{-k})_*(v_i) = \begin{cases} v_{i-n}, & \text{for } k > 0; \\ v_{i+n}, & \text{for } k < 0, \end{cases}$$

we see that

$$(4.1.7) \quad (R_{-k} \circ \varphi_{X_H})_*((\varphi_{X_H}^{-1})_*(v_i)) = \begin{cases} v_{i-n}, & \text{for } k > 0; \\ v_{i+n}, & \text{for } k < 0, \end{cases}$$

where $i, i - n$ are considered modulo $n|k|$. Similarly, since β_- and hence $d\alpha_{\tilde{M}} = d\beta_-$ is invariant under $\frac{2\pi}{n|k|}$ -rotation around p ,

$$(R_{-k})_*(w_i) = \begin{cases} w_{i-n}, & \text{for } k > 0; \\ w_{i+n}, & \text{for } k < 0. \end{cases}$$

Hence, we get

$$(4.1.8) \quad (R_{-k} \circ \varphi_{X_H})_*((\varphi_{X_H}^{-1})_*(w_i)) = \begin{cases} w_{i-n}, & \text{for } k > 0; \\ w_{i+n}, & \text{for } k < 0, \end{cases}$$

where $i, i - n$ are considered modulo $n|k|$. Formulas 4.1.7 and 4.1.8 imply that the framings given by Formula 4.1.5 and Formula 4.1.6 give rise to the symplectic trivializations $\tau(\gamma_i)$ along γ_i and $\tau(\gamma^{|k|})$ along $\gamma^{|k|}$ which are consistent in terms of Section 2.2 along the surfaces obtained from $(\varphi_{X_H}^{(-N-t)/2N}(\gamma_{p,p_i}))_{i=1}^{n|k|}$ by gluing them with φ_k . In addition, note that from the property described by Equation 4.1.4 and the fact that $d\alpha_{\tilde{M}}$ is t -invariant on V_i it follows that $w_{i,t}(p_i) = q(t)w_i(p_i)$, where $q(t) \neq 0$, $i = 1, \dots, n|k|$ and $t \in [-N, N]$. Thus, $\tau(\gamma_i)$ and τ_{γ_i} are in the same homotopy class of symplectic trivializations of the 2-plane bundle $\gamma_i^*\xi$ over S^1 . In addition, it is easy to see that the symplectic trivialization $\tau(\gamma^{|k|})$ and $\tau_{\gamma^{|k|}}$ obtained by pull-back from γ are in the same homotopy class of symplectic trivializations of the 2-plane bundle $(\gamma^{|k|})^*\xi$ over S^1 . Thus, we see that the symplectic trivializations τ_{γ_i} 's and τ_γ are consistent in the sense of Section 2.2. From now on, for simplicity we denote the consistent trivialization by τ . $\square$

Remark 4.3. If $l > 1$, one constructs τ using the method from Lemma 4.2 and get

$$\mu_\tau(\gamma_i^s) = \begin{cases} 2ls, & \text{for } k < -1; \\ -2ls, & \text{for } k > 1, \end{cases}$$

and

$$\mu_\tau(\gamma^t) = \begin{cases} 2(m-1)l + 1, & \text{for } k < -1 \text{ and } t = (m-1)|k|; \\ -2ml + 1, & \text{for } k > 1 \text{ and } t = m|k|. \end{cases}$$

for $i = 1, \dots, n$; $t \leq N_\delta$, $s \leq \frac{N_\delta}{|k|}$.

4.2. Sutured cylindrical contact homology. In this section, we calculate the sutured version of cylindrical contact homology of the sutured contact solid torus that we have constructed in Section 3.2 (in the case when $l = 1$).

We first recall that Lemma 4.1 implies that all closed orbits of $R_{\tilde{\alpha}_\delta}$ are nondegenerate.

Remark 4.4. Note that there are no contractible Reeb orbits. Hence, from Theorem 2.7, Remark 2.12, and the fact that $\pi_1(S^1 \times D^2; \mathbb{Z}) \simeq H_1(S^1 \times D^2; \mathbb{Z}) \simeq \mathbb{Z}$ it follows that for all $h \in H_1(S^1 \times D^2; \mathbb{Z})$ $HC_*^{cyl, h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta, J)$ is defined, i.e., $\partial^2 = 0$, and is independent of contact form $\tilde{\alpha}_\delta$ for the given contact structure $\tilde{\xi}$ and the almost complex structure J .

Lemma 4.2 implies that all Reeb orbits are good and

$$(4.2.1) \quad |\gamma_i^s| = \begin{cases} 2s - 1, & \text{for } k < -1; \\ -2s - 1, & \text{for } k > 1, \end{cases}$$

and

$$(4.2.2) \quad |\gamma^t| = \begin{cases} 2m - 2, & \text{for } k < -1 \text{ and } (m-1) \leq \frac{t}{|k|} < m; \\ -2m, & \text{for } k > 1 \text{ and } (m-1) < \frac{t}{|k|} \leq m \end{cases}$$

for $i = 1, \dots, n$, $s \leq \frac{N_\delta}{|k|}$ and $t \leq N_\delta$. Hence, we get

$$(4.2.3) \quad C_m^h(\tilde{\alpha}_\delta, J) = \begin{cases} \mathbb{Q}\langle \gamma^h \rangle, & \text{for } h > 0 \text{ and } m = 2[h(-\frac{1}{k} + \delta)]; \\ \mathbb{Q}\langle \gamma_1^{h/|k|}, \dots, \gamma_n^{h/|k|} \rangle, & \text{for } k \mid h > 0 \text{ and } m = -\frac{2h}{k} - 1; \\ 0, & \text{otherwise} \end{cases}$$

for $h \leq N_\delta$.

Now since by Lemma 4.1 $\mathcal{A}(\gamma^{|k|}) > \mathcal{A}(\gamma_i)$ for $i = 1, \dots, n$, we can use Fact 2.6 and Remark 2.12 and conclude that $\partial(\gamma_i^s) = 0$ for $i = 1, \dots, n$ and $s > 0$. Then we prove that $\partial(\gamma^t) = 0$ for $k \nmid t \leq N_\delta$. Since $[\gamma_i] = |k|[\gamma]$ in $H_1(S^1 \times D^2; \mathbb{Z}) \cong \mathbb{Z}$, the cylindrical contact homology differential at γ^t counts only cylinders with negative end at γ^t . Then, similarly to the previous case, Fact 2.6 and Remark 2.12 imply that $\partial(\gamma^t) = 0$ for $k \nmid t \leq N_\delta$.

We now consider the case when $k \mid t$ and will show that $\partial(\gamma^t) \neq 0$ for $k \mid t \leq N_\delta$. Is this situation, by arguing in the same way as in the case when $k \nmid t$, we get that $\partial(\gamma^t)$ counts only cylinders with negative end at $\gamma_i^{t/|k|}$.

Now we note that

$$(4.2.4) \quad \text{ind}(u) = |\gamma^t| - |\gamma_i^{t/|k|}|$$

for any pseudoholomorphic curve u in the moduli space $\mathcal{M}^J(\gamma^t; \gamma_i^{t/|k|})$, where $k \mid t \leq N_\delta$ and J is an almost complex structure tailored to $((\mathbb{R} \times S^1 \times D^2)^*, \tilde{\alpha}_\delta^*)$. The index formula can be written in this way, since $H_2(S^1 \times D^2; \mathbb{Z}) = 0$ and hence $\langle c_1(\xi), A \rangle = 0$ for all $A \in H_2(S^1 \times D^2, \mathbb{Z})$. We now use Equations 4.2.1 and 4.2.2 in the case when $k < -1$ and get

$$|\gamma^t| - |\gamma_i^{t/|k|}| = 2m - 2\frac{t}{|k|} + 1 = 2(m - \frac{t}{|k|}) + 1,$$

and $m = \frac{t}{|k|}$ for $i = 1, \dots, n$; $t \leq N_\delta$. Hence, we can rewrite Equation 4.2.4 as

$$(4.2.5) \quad \text{ind}(u) = |\gamma^t| - |\gamma_i^{t/|k|}| = 2(\frac{t}{|k|} - \frac{t}{|k|}) + 1 = 1$$

for $i = 1, \dots, n$ and $t \leq N_\delta$. Therefore, Theorem 2.8 and Remark 2.12 imply that for every $u \in \mathcal{M}(\gamma^t, \gamma_i^{t/|k|})$ the linearization of the Cauchy-Riemann operator is surjective at u ; here $k \mid t \leq N_\delta$, J is any almost complex structure tailored to $((S^1 \times D^2)^*, \tilde{\alpha}_\delta^*)$ and $i = 1, \dots, n$.

Similarly, we use Equations 4.2.1 and 4.2.2 in the case when $k > 1$ and get

$$|\gamma^t| - |\gamma_i^{t/|k|}| = -2m - (-2\frac{t}{|k|} - 1) = 2(\frac{t}{|k|} - m) + 1,$$

and $m = \frac{t}{|k|}$ for $i = 1, \dots, n$; $t \leq N_\delta$. Thus, we can rewrite Equation 4.2.4 as

$$(4.2.6) \quad \text{ind}(u) = |\gamma^t| - |\gamma_i^{t/|k|}| = 2(m - m) + 1 = 1$$

for $i = 1, \dots, n$ and $t \leq N_\delta$. Therefore, Theorem 2.8 and Remark 2.12 imply that for every $u \in \mathcal{M}(\gamma^t, \gamma_i^{t/|k|})$ the linearization of the Cauchy-Riemann operator is surjective at u ; here $k \mid t \leq N_\delta$, J is any almost complex structure tailored to $((S^1 \times D^2)^*, \tilde{\alpha}_\delta^*)$ and $i = 1, \dots, n$.

Let $(S^1 \times D^2, \Gamma_{\text{long}}, U(\Gamma_{\text{long}}), \alpha_\delta^{\text{long}})$ be a sutured contact solid torus obtained from $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$ by using φ_{X_H} as a gluing map. Recall that we get $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}), \tilde{\alpha}_\delta)$ from $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$ by using $\varphi_k = R_{-k} \circ \varphi_{X_H}$ as a gluing map. Here R_{-k} is a $-\frac{2\pi}{k}$ -rotation around the center of D . We now note that $(S^1 \times D^2, \Gamma_{\text{long}}, U(\Gamma_{\text{long}}), \alpha_\delta^{\text{long}})$ is a universally tight sutured contact solid torus with $2n|k|$ parallel longitudinal sutures, $|k| > 1$. This follows from the gluing/classification result for universally tight contact structures on a sutured solid torus, see Section 2 in [9] (more precisely, Corollary 2.3, Theorem 2.5 and Corollary 2.6). The cylindrical contact homology of this sutured contact manifold is computed in [8] and is given by

$$(4.2.7) \quad HC^{\text{cyl}, h}(S^1 \times D^2, \Gamma_{\text{long}}, \xi_{\text{long}}) \simeq \begin{cases} \mathbb{Q}^{n|k|-1}, & \text{for } h \geq 1; \\ 0, & \text{otherwise.} \end{cases}$$

Here $\xi_{\text{long}} = \ker \alpha_{\text{long}}$.

Note that $(S^1 \times D^2, \Gamma_{\text{long}}, U(\Gamma_{\text{long}}), \alpha_\delta^{\text{long}})$ has $n|k|$ hyperbolic orbits $\gamma_1^{\text{long}}, \dots, \gamma_{n|k|}^{\text{long}}$ and one elliptic orbit γ^{long} . Here γ_i^{long} 's correspond to the equally spaced saddle points of H and γ^{long} corresponds to the critical point of H at the center of $D(r_{\text{sing}})$. In addition, observe that

$$(4.2.8) \quad [\gamma_i^{\text{long}}] = [\gamma^{\text{long}}] = 1 \in H_1(S^1 \times D^2; \mathbb{Z}).$$

Finally, note that from Lemma 4.1 and from the construction of γ^{long} and $\gamma_1^{long}, \dots, \gamma_{n|k|}^{long}$ it follows that

$$(4.2.9) \quad \mathcal{A}(\gamma^{long}) > \mathcal{A}(\gamma_i^{long}), \quad \mathcal{A}(\gamma_i^{long}) = \mathcal{A}(\gamma_j^{long})$$

for $i, j = 1, \dots, n|k|$. Hence, Theorem 2.7, Remark 2.12 together with Fact 2.6, and Formulas 4.2.7, 4.2.8 and 4.2.9 imply that $\partial\gamma^{long} \neq 0$; otherwise we come to contradiction to Formula 4.2.7 ($\partial\gamma^{long} = 0$ implies that the exponent of $\mathbb{Q}$ in Formula 4.2.7 must be $n|k| + 1$).

We now take almost complex structure J^{long} tailored to $((S^1 \times D^2)^*, (\alpha_\delta^{long})^*)$ such that as a map $\xi^{long} \rightarrow \xi^{long}$ it is obtained from some fixed $J^{cyl} : \xi \rightarrow \xi$ which is defined on $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$ and satisfies the following properties:

- (1) $(J^{cyl})^2 = -I$, $d\alpha(J^{cyl}\cdot, J^{cyl}\cdot) = d\alpha(\cdot, \cdot)$, $d\alpha(\cdot, J^{cyl}\cdot) > 0$;
- (2) J^{cyl} is $\frac{2\pi}{n|k|}$ -symmetric, i.e., it is invariant under $\frac{2\pi}{n|k|}$ -rotation with respect to the center of D .

Here $\xi_{long} = \ker \alpha_{long}$ and $\xi = \ker \alpha$. By saying that J^{long} is obtained from J^{cyl} we simply mean that the gluing procedure with φ_{X_H} applied to $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$ transforms J^{cyl} to J^{long} . Since ξ is $\frac{2\pi}{n|k|}$ -symmetric on $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$, we claim that J^{cyl} , which satisfies Properties (1) and (2), exists and that Property (2) is not a serious restriction on J^{cyl} . The symmetry of ξ follows from the symmetry of β and X_H , and from the construction of α .

Now we take $\tilde{J}$ on $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}), \tilde{\alpha}_\delta)$, which is obtained from the same J^{cyl} defined on $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$ by applying the gluing procedure with $\varphi_k = R_{-k} \circ \varphi_{X_H}$ to $([-1, 1] \times D, \Gamma, U(\Gamma), \alpha)$, and possibly modify it near the boundary of $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}), \tilde{\alpha}_\delta)$ (far from the Reeb orbits) so that it becomes tailored to $((S^1 \times D^2)^*, (\tilde{\alpha}_\delta)^*)$. From the symmetry of J^{cyl} and the form of the gluing maps for $(S^1 \times D^2, \tilde{\Gamma}, U(\tilde{\Gamma}), \tilde{\alpha}_\delta)$ and $(S^1 \times D^2, \Gamma_{long}, U(\Gamma_{long}), \alpha_\delta^{long})$ we see that $\partial\gamma^{long} \neq 0$ for the count with respect to J^{long} implies that $\partial\gamma^t \neq 0$ for $k \mid t \leq N_\delta$ for the count with respect to $\tilde{J}$. Observe that this choice of almost complex structures is possible, since Theorem 2.8 and Remark 2.12 imply that we do not need to require almost complex structures to be generic. Finally, from Formula 4.2.3 it follows that

$$HC_m^{cyl, h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta) \simeq \begin{cases} \mathbb{Q}, & \text{for } h > 0 \text{ and } m = 2[h(-\frac{1}{k} + \delta)]; \\ \mathbb{Q}^{n-1}, & \text{for } k \mid h > 0 \text{ and } m = -\frac{2h}{k} - 1; \\ 0, & \text{otherwise} \end{cases}$$

for $h \leq N_\delta$.

We now note that $\tilde{\xi} = \ker \tilde{\alpha}_\delta$ is independent of δ . This follows from the gluing/classification result for universally tight contact structures on a sutured solid torus, see Section 2 in [9] (more precisely, Corollary 2.3, Theorem 2.5 and Corollary 2.6). Hence, from Theorem 2.7 and Remark 2.12 it follows that

$$HC^{cyl, h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\xi}) = HC^{cyl, h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta)$$

for all h and hence for $h \leq N_\delta$, where δ is a small positive irrational number,

$$HC^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\xi}) := \bigoplus_m HC_m^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\xi}), \quad \text{and}$$

$$HC^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta) := \bigoplus_m HC_m^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta).$$

Now observe that $N_\delta \rightarrow \infty$ when $\delta \rightarrow 0$. In addition, we note that for fixed n, k and two small positive irrational numbers $\delta_1 \neq \delta_2$, the sets of closed orbits of $R_{\tilde{\alpha}_{\delta_1}}$ and $R_{\tilde{\alpha}_{\delta_2}}$ are the same, and the corresponding orbits with the same first homology class $h \leq \min\{N_{\delta_1}, N_{\delta_2}\}$ have the same Conley-Zehnder gradings in the corresponding complexes. Therefore, for every $0 < h \in \mathbb{Z} = H_1(S^1 \times D^2; \mathbb{Z})$, there exists δ such that

$$HC_m^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\xi}) = HC_m^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\alpha}_\delta)$$

$$\simeq \begin{cases} \mathbb{Q}, & \text{for } h > 0 \text{ and } m = 2\lfloor h(-\frac{1}{k} + \delta) \rfloor; \\ \mathbb{Q}^{n-1}, & \text{for } k \mid h > 0 \text{ and } m = -\frac{2h}{k} - 1; \\ 0, & \text{otherwise} \end{cases}$$

for $h \leq N_\delta$ and hence

$$(4.2.10) \quad HC_m^{cyl,h}(S^1 \times D^2, \tilde{\Gamma}, \tilde{\xi}) \simeq \begin{cases} \mathbb{Q}, & \text{for } h > 0 \text{ and } m = 2\lfloor -\frac{h}{k} + \delta_k \rfloor; \\ \mathbb{Q}^{n-1}, & \text{for } k \mid h > 0 \text{ and } m = -\frac{2h}{k} - 1; \\ 0, & \text{otherwise,} \end{cases}$$

where $0 < \delta_k \ll \frac{1}{|k|}$. Finally, Formula 4.2.10 implies that

$$(4.2.11) \quad HC^{cyl,h}(S^1 \times D^2, \Gamma, \xi) \simeq \begin{cases} \mathbb{Q}, & \text{for } k \nmid h > 0; \\ \mathbb{Q}^{n-1}, & \text{for } k \mid h > 0; \\ 0, & \text{otherwise.} \end{cases}$$

This completes the proof of Theorem 1.1 when $l = 1$.

Remark 4.5. For $l > 1$, one can use the same observations as in the case when $l = 1$ and show that the only non-zero part of the cylindrical contact homology differential is given by $\langle \partial \gamma^t, \gamma_i^{t/|k|} \rangle \neq 0$ for $|k| \mid t \leq N_\delta$. This will lead to Formula 4.2.11 for all l such that $(k, l) = 1, |k| > l > 0$.

Remark 4.6. Theorem 1.3 from [8] and Theorem 1.1 provide the formula for the sutured version of cylindrical contact homology of $(S^1 \times D^2, \Gamma, \xi)$, where Γ is arbitrary and ξ is a universally tight contact structure. This follows from the gluing/classification result for universally tight contact structures on a sutured solid torus, see Section 2 (more precisely, Corollary 2.3, Theorem 2.5 and Corollary 2.6) in [9].

REFERENCES

- [1] F Bourgeois, *A survey of Contact Homology*, lectures at Yashafest, 2007.
- [2] F Bourgeois, Y Eliashberg, H Hofer, K Wysocki and E Zehnder, *Compactness results in symplectic field theory*, *Geom. and Top.* **7** (2003), 799–888.
- [3] V Colin, P Ghiggini, K Honda and M Hutchings, *Sutures and contact homology I*, *Geom. Topol.* **15** (2011), 1749–1842.
- [4] V Colin and K Honda, *Constructions contrôlées de champs de Reeb et applications*, *Geom. Topol.* **9** (2005), 2193–2226.

- [5] A Cotton-Clay, *Symplectic Floer homology of area-preserving surface diffeomorphisms*, Geom. Topol. **13** (2009), 2619–2674.
- [6] Y Eliashberg, A Givental and H Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal. Special Volume **10** (2000), 560–673.
- [7] D Gabai, *Foliations and the topology of 3-manifolds*, J. Diff. Geom **18** (1983), 445–503.
- [8] R Golovko, *The embedded contact homology of sutured solid tori*, Alg. Geom. Topol. **11** (2011), 1001–1031.
- [9] K Honda, *Gluing tight contact structures*, Duke Math. J. **115** (2002), 435–478.
- [10] M Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. **4** (2002), 313–361.
- [11] A Momin, *Contact Homology of orbit complements and implied existence*, preprint 2010, arXiv:1012.1386.
- [12] C Wendl, *Automatic transversality and orbifolds of punctured holomorphic curves in dimension four*, Comment. Math. Helv. **85** (2010), no. 2, 347–407.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, MONTRÉAL, QC H3T 1J4, CANADA

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, MONTRÉAL, QC H2X 3Y7, CANADA

E-mail address: `rgolovko@dms.umontreal.ca`

URL: `http://www.dms.umontreal.ca/~rgolovko`